

**PHARMACEUTICAL COMPOSITIONS COMPRISING AMLODIPINE  
MALEATE**

Patent Number: WO02053134  
Publication date: 2002-07-11  
Inventor(s): LEMMENS JACOBUS MARIA (NL); VANDERHEIJDEN ARLETTE (NL); VAN  
DALEN FRANS (NL)  
Applicant(s): LEMMENS JACOBUS MARIA (NL); VANDERHEIJDEN ARLETTE (NL); BIOORG  
B V (NL); VAN DALEN FRANS (NL)  
Requested  
Patent: ☐ WO02053134  
Application  
Number: WO2001NL00606 20010815  
Priority Number  
(s): US20000258562P 20001229; US20010809346 20010316  
IPC  
Classification: A61K9/20; A61K9/48; A61K31/44  
EC Classification: A61K31/44R  
Equivalents: ☐ DE20116428U  
Cited Documents: WO9534299; WO9505822; WO0174390; EP1181932

**Abstract**

An amlodipine maleate pharmaceutical composition is provided with good stability when formulated with a pH within the range of 5.5 to 7, when measured as a 20 wt% aqueous slurry. The stability can also be aided by making the pharmaceutical composition from amlodipine maleate particles having an average particle size of greater than 20 microns, preferably greater than 100 microns.

Data supplied from the esp@cenet database - I2

Herrn Dr. Tiedeman - Stephen Jones

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number  
WO 02/053134 A1

(51) International Patent Classification: A61K 9/20,  
9/48, 31/44

(21) International Application Number: PCT/NL01/00606

(22) International Filing Date: 15 August 2001 (15.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/258,562 29 December 2000 (29.12.2000) US  
09/809,346 16 March 2001 (16.03.2001) US

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CY, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): BIOORGANICS B.V. (NL/NL); Microweg 22, NL-6545 CM Nijmegen (NL).

(72) Inventors; and

(73) Inventors/Applicants (for US only): LEMMENS, Jacobus, Maria (NL/NL); Lodewijkstraat 30, NL-6585 KM Mook (NL). VAN DALEN, Frans (NL/NL); Groenbeekweg 11, NL-6524 CK Nijmegen (NL). VAN DERHEIJDEN, Arlette (NL/NL); Gmiederswaarsstraat 69, NL-6512 ER Nijmegen (NL).

(74) Agent: DUXBURY, Stephen; Arnold & Siedsma, Sweet-Inckpoin 1, NL-2517 GK The Hague (NL).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

= DE 201 16 428 U

WO 02/053134 A1

(54) Title: PHARMACEUTICAL COMPOSITIONS COMPRISING AMLODIPINE MALEATE

(57) Abstract: An amlodipine maleate pharmaceutical composition is provided with good stability when formulated with a pH within the range of 5.5 to 7, when measured as a 20 wt% aqueous slurry. The stability can also be aided by making the pharmaceutical composition from amlodipine maleate particles having an average particle size of greater than 20 microns, preferably greater than 100 microns.

WO 02/053134


PCT/NL01/00606

## PHARMACEUTICAL COMPOSITIONS COMPRISING AMLODIPINE MALEATE

5

The present invention relates to pharmaceutical compositions comprising amlodipine maleate and to processes for making the same.

Calcium channel blockers are useful in treating a variety of cardiac conditions, primarily angina and hypertension. EP 089 167 and corresponding US 4,572,909  
10 disclose a class of substituted dihydropyridine derivatives as being useful calcium channel blockers. These patents identify that one of the most preferred compounds is 2-[(2-aminoethoxy)methyl]-4-(2-chlorophenyl)-3-ethoxycarbonyl-5-methoxycarbonyl-6-methyl-1,4-dihydropyridine. This compound, which is now commonly known as amlodipine, has the following formula:



15

Examples 8, 11, 12, and 22 of EP 089 167 show the synthesis of amlodipine in the maleate salt form. While a variety of acid addition salts are taught to be suitable, the maleate salt is identified as the most preferred acid addition salt.

Subsequently, EP 244 944 and corresponding US 4,879,303 were issued  
20 directed to the besylate (or benzene sulfonate) salt of amlodipine. The besylate salt is stated to provide certain advantages over the known salts including good formulating properties. Indeed, amlodipine besylate, and not amlodipine maleate, has been developed into a commercial prescription pharmaceutical by Pfizer and is sold in the U.S. under the trade name NORVASC.

25

A review of the available portions of the NORVASC (amlodipine besylate) New Drug Application (NDA) filed with the U.S. Food & Drug Administration by Pfizer reveals that a switch was made during development from the original amlodipine maleate to the amlodipine besylate. (See "Review of Original NDA" for

WO 02/053134

PCT/NL01/00606

2

NDA# 19-787 of 10.10.1990, obtainable from FDA under Freedom of Information Act ). The reasons for the switch were apparently tableting and stability problems. However, the precise stability and tableting problems/issues/causes are not publicly disclosed in the information available from the FDA. Interestingly, clinical studies in  
5 the NDA include the use of the maleate salt form and the besylate salt form, the two salt forms being therapeutically equivalent (bioequivalent). However, in these studies amlodipine maleate was always used in a capsule or solution dosage form, not a tablet dosage form.

The present invention relates to the discovery of stable amlodipine maleate  
10 pharmaceutical compositions. A first aspect of the invention relates to a pharmaceutical composition comprising an effective amount of amlodipine maleate and at least one pharmaceutically acceptable excipient wherein the composition has a pH within the range of 5.5 - 7.0. The composition is preferably a solid dosage form such as a tablet or capsule. The invention also relates to methods of making such a  
15 composition as well as to the use of such compositions in treating or preventing angina or hypertension.


Another aspect of the invention relates to a process which comprises mixing solid particles of amlodipine maleate having an average particle size of at least 20 microns with one or more pharmaceutically acceptable excipients to form a mixture.  
20 Preferably the mixture has a pH of from 5.5 to 7.0. The mixture can be compressed into tablets or filled into capsules to form a solid dosage form.

Stability is an important aspect of a pharmaceutical composition. The present invention is based on the discovery that the prior art stability problems associated with amlodipine maleate can be overcome primarily by controlling the pH of the  
25 composition to be within the range of about 5.5 to 7.0, preferably of about 6.0-7.0. Within this range the potential degradation reactions are minimized. In particular, the formation of the following degradation product, referred to herein as amlodipine aspartate, is diminished or prevented in this pH range.

WO 02/053134


PCT/NL01/00606

3

**Amlodipine Aspartate**

5 Amlodipine aspartate is formed by a Michael addition reaction between amlodipine and maleic acid. Because amlodipine and maleic acid are in intimate contact with each other in the amlodipine maleate salt, the chances of the addition reaction occurring increase with time; hence raising a stability issue. By controlling the pH, the addition is significantly slowed or prevented altogether. Thus, it has now  
10 been discovered that pH levels above 7.0 tend to encourage the degradation of amlodipine maleate into amlodipine aspartate. Below a pH of about 5.5, other degradation reactions tend to be encouraged, including the pyridine analogue of amlodipine having the following structure:

15 "Amlo-pyridine"



20 Preferably the pH of the stabilized composition of our invention is in the range of about 6.0 to 7.0 and more typically from about 6.1 or 6.2 to 6.8. For solid compositions, the pH is determined by forming a slurry of the material with water (demineralized water) and measuring the pH of the slurry, as is understood by workers skilled in the art regarding the pH of a solid composition. The concentration of the composition in the slurry is 20 wt%. The pH is measured by any standard technique.

The pharmaceutical compositions of the present invention comprise a pharmaceutically effective amount of amlodipine maleate and at least one

WO 02/053134

PCT/NL01/00606

4

pharmaceutically acceptable excipient. Preferably the stability of the composition is such that after three months, more preferably after six months, in an environment controlled room at 40°C/75% RH, it exhibits a loss of amlodipine (or, accordingly, an increase in impurities content) of less than 10 %, preferably less than 5 %, and more preferably less than 1%. Alternatively, the amlodipine maleate pharmaceutical compositions of the present invention preferably exhibit a storage stability equivalent to or superior to amlodipine besylate compositions. For example, the loss of amlodipine during storage to degradation reactions is equivalent to (+/- 10%) or less than the amlodipine loss in amlodipine besylate compositions, especially the commercial product.

The form of the amlodipine maleate is not particularly limited and includes anhydrides, solvates, hydrates and partial hydrates as well as crystalline and amorphous forms. Further, the ratio of amlodipine to maleate can be varied and specifically includes the more common and prior art form of 1:1 as well as a novel 2:1 form described commonly owned co-pending U.S. patent application serial No. 09/809,356, filed on March 16, 2001, and entitled "Amlodipine Hemimaleate," the entire contents of which are incorporated herein by reference.

The amount of amlodipine is not particularly limited and includes any amount that provides a pharmaceutical effect. In particular, amlodipine maleate can be used to treat or prevent hypertension or angina by administering an effective amount to a patient in need thereof. The specific form of angina is not particularly limited and specifically includes chronic stable angina pectoris and vasospastic angina (Prinzmetal's angina). The compound can be administered by any suitable route including orally or parenterally depending on the dosage form. The "patients" intended to be treated include human and non-human animals especially non-human mammals. Generally the amount of amlodipine maleate in a unit dose is from 1 to 100 mg, more typically from 1 to 25 mg, and preferably about 1, 1.25, 2.5, 5 or 10 mg (expressed in terms of the free base).

Amlodipine maleate can be made by any of the known techniques set forth in the prior art, including those described in the above-mentioned patents EP 089 167 and US 4,572,909. It is desirable that the amlodipine maleate active be substantially pure. For example, the content of impurities such as amlodipine aspartate, amlodipine, etc. that can be produced during the synthesis should be limited, preferably to less than 2 wt %, although such purity is not required for the present invention. A

WO 02/053134

PCT/NL01/00606

5

useful method to produce amlodipine maleate substantially free from amlodipine aspartate is described in commonly owned co-pending U.S. patent application serial No. 09/809,343, filed on March 16, 2001, and entitled "Process for Making Amlodipine Maleate," the entire contents of which are incorporated by reference.

- 5 Similarly, a useful method for making amlodipine free base is described in commonly owned co-pending U.S. patent application serial No. 09/809,351, filed on March 16, 2001, and entitled "Process For Making Amlodipine, Derivatives Thereof, and Precursors Therefor," the entire contents of which are incorporated herein by reference.

- 10 The pharmaceutical compositions of the present invention also contain at least one excipient. An "excipient" as used herein means any pharmaceutically acceptable inactive component of the composition. As is well known in the art, excipients include diluents, binders, lubricants, disintegrants, colorants, preservatives, pH-adjusters etc. The excipients are selected based on the desired physical aspects of the
- 15 final form: e.g., obtaining a tablet with desired hardness and friability, being rapidly dispersible and easily swallowed, etc. The desired release rate of the active substance from the composition after its ingestion also plays a role in the choice of excipients. Preferred release rate is the rate comparable with commercially available amlodipine besylate tablets.

- 20 Suitable excipients for use in this invention include:

- a diluent such as calcium hydrogen phosphate, lactose, mannitol etc.
- a binder such as microcrystalline cellulose or a modified cellulose, povidone etc.
- a disintegrant such as sodium starch glycolate, croscopolvidone
- 25 - a lubricant such as magnesium stearate, sodium stearyl fumarate, talc
- a colorant, taste masking agent etc.,

- The pH of the composition can be controlled or adjusted by the proper selection of excipients. It should be borne in mind that amlodipine maleate is slightly acidic. For example, amlodipine maleate has a pH of about 4.8 as a saturated aqueous
- 30 solution. Thus using excipients that are pH inert; i.e., they have little to no effect on pH, generally results in a non-alkaline pharmaceutical composition because the amlodipine maleate essentially acts as its own adjusting agent. An example of a pH-inert excipient is microcrystalline cellulose. A composition comprising amlodipine maleate and microcrystalline cellulose generally exhibits a pH of about 6. For

WO 02/053134

6

PCT/NL01/00606

comparison, the corresponding amlodipine besylate composition generally has a pH of about 7 and the corresponding amlodipine free base composition generally has a pH of about 9. Commercially available tablets comprising amlodipine besylate and sold under brand name Norvasc exhibit a pH typically between 7.05-7.35 (again measured as a 20 wt% slurry).

Excipients having a pH effect can also be used. The pH of these excipients must be taken into account in developing the pharmaceutical composition so that the overall pH of the pharmaceutical composition falls within the range of about 5.5 to 7.0.

For example, the commercially available/pharmaceutically acceptable calcium phosphates are generally alkaline; i.e. pH greater than 7 when measured as described above in a 20% slurry. For instance, DI-TAB, a commercially available dibasic calcium phosphate dihydrate, is reported as having a pH of about 7.4. Nonetheless some forms and grades of calcium phosphate are acidic or neutral pH. This lower pH can be due to the species of calcium phosphate as well as the treatment during processing of the material, such as in removing impurities/washing. For example, dibasic calcium phosphate anhydrate is generally considered to have a pH of about 7.3 whereas A-Tab™ (Rhodia), also a dibasic calcium phosphate anhydrate, has a pH of about 5.0. Further examples of commercially available non-alkaline calcium phosphates include DiCAFOS A (Budenheim) having a pH of about 7 (10 % slurry) and Fujicalin SG (Fuji) having a pH from 6.1-7.2 (5 % slurry). By using a non-alkaline calcium phosphate as an excipient, a pharmaceutical composition meeting the desired pH can be attained. Alternatively, a blend of calcium phosphates, some at pH above 7 and some a pH below 7 can be used to achieve the desired pH of the composition.

Instead of, or in addition to a non-alkaline calcium phosphate, other acidic excipients can be used by themselves or to counter balance an alkaline excipient. An example of such an acidic excipient is the disintegrant Explotab(TM) of Penwest, which is a cross-linked, low substituted sodium starch glycolate. Further, pH adjusting agents can also be used to attain the desired pH. These agents include pharmaceutically acceptable acids such as maleic acid, citric acid or ascorbic acid (the second two have utility also to act as antioxidants) and pharmaceutically acceptable bases such as calcium oxide or magnesium oxide. Salts of weak acids and/or weak



WO 02/053134

PCT/NL01/00606

7

bases are also suitable pH adjusters as they act as buffers imparting pH to lower or higher values according to the chemical nature of their components.

The pharmaceutical compositions of the present invention are not particularly limited in terms of form or route of administration. Oral dosage forms as well as parenteral dosage forms are included. The composition can be in the form of a liquid, solid, or suspension. Preferably, the pharmaceutical composition is a solid dosage form such as a tablet, capsule or sachet intended for oral administration.

Preferred solid dosage forms contain as a major excipient microcrystalline cellulose, a calcium phosphate, especially calcium hydrogenphosphate, or a combination thereof. The sum of the other excipients, if any, is generally less than 25 wt%, more usually less than 10 wt% and in some cases less than 5 wt% of the total pharmaceutical composition. Preferred other excipients are a disintegrant such as sodium starch glycollate and/or a lubricant such as magnesium stearate and/or talc.

For example, a pharmaceutical composition comprising amlodipine maleate and microcrystalline cellulose as the only excipient has been shown to provide good stability against the formation of impurities.

The pharmaceutical compositions of the present invention can be made by techniques generally known in the art. In general, the amlodipine maleate is mixed with one or more excipients to form a blend. The mixing can be carried out wet or dry (i.e. using or not using a solvent or a liquid diluent in the process) and can involve granulating, slugging, or blending of powders. A dry process is however preferred. The blend, after optional further processing, can be compressed into tablets or filled into capsules such as gelatin capsules. Typically the amlodipine maleate to be mixed is in the form of particles. The storage stability of the pharmaceutical composition of the present invention is enhanced, in general, by using larger particle sizes. Preferably the average particle size of the amlodipine maleate is at least 20 microns, more preferably at least 100 microns, and in some embodiments at least 300 microns. If an adjustment of the pH of the composition is needed, it preferably is adjusted before processing into the final form such as a tablet or capsule.

For example, tablets made of the present invention can be prepared, e.g. by a wet granulation of a mixture of amlodipine maleate and a solid carrier/diluent such as calcium phosphate of the proper grade, with the aid of a granulating solvent such as water or ethanol; drying the wet granulate; sieving the granulate; blending with sodium starch glycollate and magnesium stearate and compressing the tablet mixture

WO 02/053134

PCT/NL01/00606

8

For example, tablets made of the present invention can be prepared, e.g. by a wet granulation of a mixture of amlodipine maleate and a solid carrier/diluent such as calcium phosphate of the proper grade, with the aid of a granulating solvent such as water or ethanol; drying the wet granulate; sieving the granulate; blending with sodium starch glycollate and magnesium stearate and compressing the tablet mixture into tablets. The control of pH and/or adjusting the pH value should be advantageously included before blending with magnesium stearate. In this example both the granulating step and the blending step are considered as "mixing" steps since amlodipine maleate and an excipient are mixed.

Yet another suitable process comprises direct compression of the blend of amlodipine maleate and excipient(s). In this process, the ingredients are blended together to form a compressible blend composition that is subsequently compressed into a tablet. A blend that comprises amlodipine maleate, microcrystalline cellulose and/or calcium phosphate, and optionally sodium starch glycollate and/or magnesium stearate may be useful in forming a tablet by direct compression. For example, a blend comprising amlodipine maleate, calcium hydrogen phosphate, microcrystalline cellulose and sodium starch glycollate having a pH of 5.5 to 7.0 can be blended together, re-blended with magnesium stearate and pressurized to form a stable tablet.

The tablets of the present invention comprising amlodipine maleate, microcrystalline cellulose, sodium starch glycollate and magnesium stearate, optionally with addition of calcium hydrogenphosphate, does not suffer from the problem of stickiness to the tablet punch as reported in prior art regarding other amlodipine formulations (see the above cited EP 244 944). Accordingly, the composition of the present invention may be produced in industrial scale without technological problems.

Tablets may be covered with a suitable coating. For example, the coating can be a moisture barrier to help with storage stability or a sustained or delayed release coating composition as are well known in the art.

Alternative dosage forms are capsules, both soft and hard capsules. The stabilized amlodipine maleate composition of the invention as described above is filled into capsules by techniques known in the art, in amounts comprising the desired therapeutical dose of amlodipine.

WO 02/053134

PCT/NL01/00606

9

Suitable package material for packing the pharmaceutical dosage forms are plastic or glass containers and blisters. Particularly blisters made from non-permeable materials (high density polyethylene or aluminium) are advantageous as they may contribute to decreasing the rate of formation of degradation impurities, namely the amlo-pyridine impurity, during storage.

The pharmaceutical compositions of the present invention are used in treating or preventing angina or hypertension, as previously defined, by administering an effective amount of the pharmaceutical composition to a patient in need thereof. Typically the pharmaceutical composition is a unit dose. Individual unit dose compositions generally contain from 1 to 100 mg of amlodipine maleate, more usually from 1 to 25 mg. Preferable are unit doses comprising amlodipine maleate in an equivalent of 1.25, 2.5, 5 or 10 mg of amlodipine, such as tablets or capsules for oral administration. The pharmaceutical composition is administered from 1 to 3 times daily, preferably once a day. The above compositions are also useful in reducing heart failure symptoms, improving systolic left ventricular function and increasing exercise capacity in patients with ischaemic LVD and heart failure without current angina.

Amlodipine maleate compositions of our invention may be also used in medical applications in combination with other antihypertensive and/or antianginal agents, for instance with ACE- inhibitors such as benazepril. The combination may be realized in a form of single combination preparation, e.g. a capsule containing amlodipine maleate and benazepril hydrochloride, or by separate administration of drugs containing the above agents. Similarly, amlodipine maleate may also be combined with HMG-CoA reductase inhibitors, particularly statins such as lovastatin, simvastatin, atorvastatin, etc.

Accordingly, the present invention further provides a method for treating and/or preventing any one or more of angina, hypertension and heart failure (the Disorders) by administering a pharmaceutical composition of our invention comprising an effective and/or prophylactic amount of amlodipine maleate to a sufferer in need thereof.

The present invention also provides the use of the composition of the invention in the manufacture of a medicament for treating and/or preventing any one or more of the Disorders.

WO 02/053134

10

PCT/NL01/00606

**Examples****EXAMPLE 1 Amlodipine maleate tablets based on calcium phosphate excipient**

5

**a) Tablet compositions comprising calcium phosphate of different pH**

Batch number	(A)	(B)	(C)	(D)	(E)	(F)
Equivalent of amlodipine	2.5 mg	10 mg	2.5 mg	10 mg	2.5 mg	10 mg
Amlodipine maleate	3.21 mg	12.8 mg	3.21 mg	12.8 mg	3.21 mg	12.8 mg
Calcium hydrogen phosphate anhydrous: - Di CAFOS A - A-TAB - Fujicalin	31.5 mg	126.0 mg	31.5 mg	126.0 mg	31.5 mg	126.0 mg
Microcrystalline cellulose	62.05 mg	248.1 mg	62.05 mg	248.1 mg	62.05 mg	248.1 mg
Sodium starch Glycollate	2.0 mg	8.0 mg	2.0 mg	8.0 mg	2.0 mg	8.0 mg
Magnesium Stearate	1.0 mg	4.0 mg	1.0 mg	4.0 mg	1.0 mg	4.0 mg
Total	99.76 mg	398.9 mg	99.76 mg	398.9 mg	99.76 mg	398.9 mg

**Types of anhydrous calcium hydrogenphosphate used**

10

Batch no.	Final pH of 20% (m/V) slurry of tabs	Type of $\text{CaHPO}_4$	Supplier of $\text{CaHPO}_4$	pH of 5% (m/V) slurry of $\text{CaHPO}_4$	pH of 20% (m/V) slurry of $\text{CaHPO}_4$
(A), (B)	6.13, 6.19	Di CAFOS A	Budenheim	7.29	6.69
(E), (F)	5.74, 5.74	Fujicalin	Fuji	6.12	5.62
(C), (D)	5.53, 5.54	A-TAB	Rhone-Poulenc	6.03	5.25

WO 02/053134

PCT/NL01/00606

11

## b) Tablet compositions comprising amlodipine maleate of different particle sizes

Batch number	(G)	(H)	(I)
Equivalent of amlodipine base	2.5 mg	10 mg	2.5 mg
Amlodipine maleate, milled	3.21 mg	12.8 mg	-
Amlodipine maleate	-	-	3.21 mg
Calciumhydrogenphosphate anhydrous (pH 6.7)	31.5 mg	126.0 mg	31.5 mg
Microcrystalline cellulose	62.05 mg	248.1 mg	62.05 mg
Sodium starch glycolate	2.0 mg	8.0 mg	2.0 mg
Magnesium Stearate	1.0 mg	4.0 mg	1.0 mg
Total	99.76 mg	398.9 mg	99.76 mg

- 5 Particle size of the amlodipine maleate substance used for production of batches A – F was measured by laser diffraction and it was proven that less than 90% of particles is smaller than 204 microns and 50% of particles is smaller than 80 microns.

- 10 This amlodipine maleate was milled to a particle size of 10-20 microns and used for producing batches (G) and (H). Alternately, another batch of amlodipine maleate with particle sizes 90% less than 11 microns and 50% less than 6 microns was used for producing batch (I).

## c) Manufacturing procedures

- 15 Batches (A) – (F) and (I) have been manufactured as follows:

- The amlodipine maleate was sieved through a 500 µm screen,
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- 20 - Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg and / or 10 mg tablets have been compressed using a Korsch EKO excenter press.

WO 02/053134

12

PCT/NL01/00606

Batches (G) and (H) have been manufactured as follows:

- Amlodipine maleate was milled
- The other excipients have been sieved through a 850 µm screen.
- 5 - All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg / 10 mg tablets have been compressed on Korsch EKO excenter press.

10

Example 2 Tablets comprising microcrystalline cellulose

a) Composition

Batch number	(J)	(K)
Equivalent of amlodipine base	2.5 mg	10 mg
Amlodipine maleate	3.21 mg	12.8 mg
Microcrystalline cellulose	75.55 mg	302.1 mg
Predried potato starch	20.0 mg	80.0 mg
Magnesium Stearate	0.5 mg	2.0 mg
Talc	0.5 mg	2.0 mg
Total	99.76 mg	398.9 mg

15

Manufacturing process

- Amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate and talc have been mixed in  
20 a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate and talc were added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg and 10 mg tablets have been compressed using a Korsch EKO  
25 excenter press.

WO 02/053134

13

PCT/NL01/00606

Properties of tablet composition :

pH of 20% (w/V) slurry : - Batch (J) = 5.92

- Batch (K) = 5.96

5

EXAMPLE 3 Amlodipine maleate tablets comprising mannitol

Composition

Batch number	(L)
Equivalent of amlodipine base	10 mg
Amlodipine maleate	12.8 mg
Mannitol	370.1 mg
Sodium starch glycollate	8.0 mg
Magnesium Stearate	6.0 mg
Talc	2.0 mg
Total	398.9 mg

10

Manufacturing procedure

- Amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate and talc have been mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.

15

- Magnesium stearate and talc were added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 10 mg tablets have been compressed using a Korsch EKO excenter press.

WO 02/053134

14

PCT/NL01/00606

**Example 4 Amlodipine maleate tablets having pH higher than 7**  
**(comparative example)**

Composition (pH of the 20% w/v slurry : 8.68 )

Batch number	(X)
Equivalent of amlodipine base	2.5 mg
Amlodipine maleate	3.21 mg
Calcium hydrogenphosphate anhydrous (pH 6.69)	31.5 mg
Magnesium oxide ponderosum	0.5 mg
Microcrystalline cellulose	62.05 mg
Sodium starch glycolate	2.0 mg
Magnesium Stearate	1.0 mg
Total	100.26 mg

**Manufacturing process**

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- Amlodipine maleate, magnesium oxide and about 30 % of the amount of microcrystalline cellulose (MCC) were mixed in a free fall mixer for 10 minutes at about 25 rpm.
- The remaining amount of MCC, calcium hydrogenphosphate anhydrous and sodium starch glycolate were added and the blend was mixed in a free fall mixer for 15 minutes at about 25 rpm. Value of pH was checked at 20% aqueous slurry.
- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- 2.5 mg tablets and proportionately larger 10 mg tablets have been compressed using a Korsch EK0 excenter press.



WO 02/053134

15

PCT/NL01/00606

Example 5 Stability studies on amlodipine maleate tablets

Stability studies on batches produced in Examples 1-4 were performed in a thermostated chamber adjusted to  $40 \pm 2$  °C and  $75 \pm 5$  % of relative humidity in various package materials ( HDPE bottles, PVC/PVDC/PE blisters ) or on an open dish. Assay of the active substance and of the content of impurities was performed by HPLC method, using reference materials of amlodipine maleate and major degradation impurities :

amlodipine aspartate ( Z#204)

amlo-pyridine (Z# 202)

10 Apart, two minor impurities Z#203 and Z#205 were detected and identified.

The content of other detected impurities/degradation products was calculated by internal normalization.

In the following tables, the assay of the active substance is expressed in milligrams, the content of impurities is expressed in per cent.

WO 02/053134

16

PCT/NL01/00606

A) Stability studies performed at 40°C/75%RH, open dish (influence f pH)

t=0 months	(A)	(B)	(E)	(F)	(C)	(D)
Assay (mg/tab)	2.45	9.85	2.38	10.01	2.38	9.88
Z#202 (%)	0.15	0.15	0.08	0.08	0.05	0.08
Z#203 (%)	0.00	0.03	0.03	0.03	0.03	0.03
Z#204 (%)	0.16	0.15	0.13	0.12	0.13	0.12
Z#205 (%)	0.01	0.01	0.00	0.00	0.00	0.00
Tot.unknown(%)	0.31	0.28	0.27	0.28	0.27	0.27
T= 1 month						
Assay (mg/tab)	2.48	9.82	2.31	9.55	2.23	9.29
Z#202 (%)	0.15	0.15	0.21	0.23	1.27	1.25
Z#203 (%)	0.00	0.03	0.04	0.04	0.04	0.04
Z#204 (%)	0.24	0.23	0.41	0.30	0.27	0.22
Z#205 (%)	0.01	0.01	0.03	0.02	0.04	0.03
Tot.unknown(%)	0.35	0.31	0.58	0.56	0.83	0.72
T= 3 months						
Assay (mg/tab)	2.40	9.69	2.30	9.57	2.16	8.94
Z#202 (%)	0.20	0.18	0.37	0.39	1.85	1.85
Z#203 (%)	0.03	0.03	0.04	0.05	0.04	0.04
Z#204 (%)	0.31	0.29	0.77	0.63	0.33	0.27
Z#205 (%)	0.01	0.01	0.05	0.04	0.03	0.02
Tot. Unkn. (%)	0.42	0.35	1.11	1.10	1.37	1.10

WO 02/053134

17

PCT/NL01/00606

Stability studies performed in PVC/PE/PVDC blister at 40°C/75%RH

T= 0 months	(A)	(B)	(J)	(K)
Assay	2.51	9.99	2.52	10.33
Z#202	0.11	0.11	0.20	0.20
Z#203	0.00	0.00	0.00	0.00
Z#204	0.15	0.14	0.00	0.00
Z#205	0.01	0.01	0.00	0.00
T unkn.	0.31	0.31	0.46	0.49
T= 3 months				
Assay	2.49	9.49	2.49	9.99
Z#202	0.15	0.15	0.07	0.05
Z#203	0.00	0.00	0.00	0.03
Z#204	0.26	0.29	0.46	0.40
Z#205	0.01	0.01	0.02	0.02
T unkn.	0.36	0.34	0.45	0.38
T= 6 months				
Assay	2.45	9.49	2.45	9.94
Z#202	0.23	0.21	0.17	0.11
Z#203	0.04	0.04	0.04	0.04
Z#204	0.46	0.39	0.64	0.54
Z#205	0.01	0.00		0.02
T unkn.	0.57	0.54	0.64	0.51

5 Stability studies performed open dish at 40°C/75%RH

t=0 months	(J)	(K)
Assay	2.49	10.18
Z#202	0.04	0.04
Z#203	0.02	0.03
Z#204	0.19	0.17
Z#205	0.01	0.01
tot unkn.	0.42	0.36
t=1 month		
Assay	2.44	10.14
Z#202	0.08	0.07
Z#203	0.02	0.03
Z#204	0.34	0.36
Z#205	0.01	0.01
tot unkn.	0.45	0.42
t=3 months		
assay	2.41	9.72
Z#202	0.22	0.21
Z#203	0.04	0.04
Z#204	0.50	0.54
Z#205	0.03	0.03
tot unkn.	0.63	0.56

WO 02/053134

18

PCT/NL01/00606

Stability studies performed in HDPE container at 40°C/75%RH

T= 0 months	(A)	(B)	(J)	(K)	(L)
Assay	2.51	9.99	2.52	10.33	10.11
Z#202	0.11	0.11	0.20*	0.20*	0.19*
Z#203	0.00	0.00	0.00	0.00	0.00
Z#204	0.15	0.14	0.00*	0.00*	0.00*
Z#205	0.01	0.01	0.00	0.00	0.00
T unkn.	0.31	0.31	0.46	0.49	0.40
T= 3 month					
Assay	2.52	9.84	2.51	10.26	8.69
Z#202	0.13	0.15	0.41*	0.37*	0.45*
Z#203	0.00	0.00	0.03	0.00	0.05
Z#204	0.27	0.23	0.00*	0.00*	0.00*
Z#205	0.03	0.03	0.07	0.05	0.01
T unkn.	0.33	0.32	0.41	0.49	0.66
T= 6 months					
Assay	2.49	9.83	2.47	10.26	
Z#202	0.15	0.15	0.08	0.05	
Z#203	0.00	0.00	0.00	0.03	
Z#204	0.44	0.41	0.44	0.38	
Z#205	0.02	0.01	0.10	0.08	
T unkn.	0.46	0.44	0.44	0.37	

\* Values for Z#202 and Z#204 together

- 5 Comparative stability studies performed at the composition with alkaline pH (open dish, 40°C/75%RH

t=0 months	(X)
Assay	2.52
Z#202	0.04
Z#203	0.03
Z#204	0.13
Z#205	0.00
tot unkn.	0.30
t=1 month	
Assay	2.43
Z#202	0.06
Z#203	0.03
Z#204	1.73
Z#205	0.00
tot unkn.	0.51

WO 02/053134

19

PCT/NL01/00606

Stability studies performed at 40°C/75%RH, open dish (influence of particle size)

t = 0 months	(A)	(B)	(G)	(H)	(I)
Assay (mg/tab)	2.45	9.85	2.38	10.12	2.54
Z#202 (%)	0.15	0.15	0.05	0.05	0.10
Z#203 (%)	0.00	0.03	0.00	0.01	0.04
Z#204 (%)	0.16	0.15	0.03	0.03	0.01
Z#205 (%)	0.01	0.01	0.00	0.00	0.02
Tot unkn. (%)	0.31	0.28	0.17	0.11	0.26
t = 1 month					
Assay (mg/tab)	2.48	9.82	2.29	9.88	-
Z#202 (%)	0.15	0.15	0.17	0.14	-
Z#203 (%)	0.00	0.03	0.01	0.01	-
Z#204 (%)	0.24	0.23	0.52	0.48	-
Z#205 (%)	0.01	0.01	0.02	0.02	-
Tot unkn. (%)	0.35	0.31	0.22	0.19	-
t = 2 months					
Assay (mg/tab)	2.46	9.87	2.22	9.64	2.35
Z#202 (%)	0.17	0.16	0.28	0.25	0.34
Z#203 (%)	0.00	0.00	0.01	0.01	0.03
Z#204 (%)	0.28	0.28	1.02	1.01	0.41
Z#205 (%)	0.01	0.01	0.03	0.03	0.04
Tot unkn. (%)	0.47	0.43	0.63	0.52	0.60
t = 3 months					
Assay (mg/tab)	2.40	9.69	2.24	9.36	2.36
Z#202 (%)	0.20	0.18	0.18	0.16	0.39
Z#203 (%)	0.03	0.03	0.01	0.01	0.11
Z#204 (%)	0.31	0.29	1.26	1.34	0.53
Z#205 (%)	0.01	0.01	0.05	0.04	0.04
Tot unkn. (%)	0.42	0.35	0.90	0.74	0.64

- 3 For a comparison of stability, increase in total impurities, identified and unidentified, with regard to the t=0 months level, is considered:

Increase in total impurities in %	(A)	(B)	(G)	(H)	(I)
After 3 months at 40°C/75%RH	+0.33	+0.24	+1.82	+1.43	

WO 02/053134

20

PCT/NL01/00606

Comparative stability study with Norvasc (commercial amlodipine besylate tablets)

Stability studies performed in original blister at 40°C/75%RH

5

		Norvasc® 5 mg batch 81040100 (DE)	Norvasc® 10 mg batch 901-05941 (NL)
t = 0 months			
Assay		5.19	9.99
Z#202		0.04	0.02
Z#203		0.01	0.00
Z#204		0.00	0.00
Z#205		0.00	0.00
tot. unkn.		0.12	0.12
t = 3 months			
Assay		5.13	9.70
Z#202		0.16	0.17
Z#203		0.03	0.00
Z#204		0.00	0.00
Z#205		0.00	0.00
tot. unkn.		0.38	0.62
t = 6 months			
Assay		4.97	9.58
Z#202		0.28	0.27
Z#203		0.00	0.00
Z#204		0.00	0.00
Z#205		0.00	0.00
tot. unkn.		0.49	0.78

WO 02/053134

21

PCT/NL01/00606

Stability studies performed on open dish at 40°C/75%RH

		Norvasc® 2.5 mg batch 8QP115A (US)	Norvasc® 10 mg batch N-09 (ES)
t = 0 months	Assay	2.44	9.91
	Z#202	0.08	0.82
	Z#203	0.00	0.00
	Z#204	0.00	0.00
	Z#205	0.00	0.00
	tot. unkn.	0.02	0.34
t = 1 months	Assay	2.44	8.90
	Z#202	0.18	2.17
	Z#203	0.04	0.19
	Z#204	0.00	0.01
	Z#205	0.00	0.00
	tot. unkn.	0.10	1.21
t = 2 months	Assay	2.39	7.98
	Z#202	0.27	3.24
	Z#203	0.00	0.00
	Z#204	0.00	0.00
	Z#205	0.00	0.03
	tot. unkn.	0.33	2.51
t = 3 months	Assay	2.34	7.68
	Z#202	0.37	3.98
	Z#203	0.00	0.00
	Z#204	0.00	0.00
	Z#205	0.00	0.03
	tot. unkn.	0.27	2.76

5

#### Example 6 Amlodipine maleate capsules

#### Compositions

10

Batch number	(CA), (CB)
Equivalent to amlodipine	5.0 mg
Amlodipine maleate	6.42 mg
Microcrystalline cellulose	72.6 mg
Predried potato starch	20.0 mg
Magnesium stearate	0.5 mg
Total	99.52 mg

WO 02/053134

22

PCT/NL01/00606

Batch number	(CC)	(CX)
Equivalent to amlodipine	5.0 mg	5.0 mg
Amlodipine maleate	6.42 mg	6.42 mg
Calcium hydrogenphosphate anhydrous	31.5 mg	31.5 mg
Magnesium oxide	-	1.5 mg
Microcrystalline cellulose	62.0 mg	62.0 mg
Sodium starch glycolate	2.0 mg	2.0 mg
Magnesium stearate	1.0 mg	1.0 mg
Total	102.92 mg	104.43 mg

pH of capsules at t=0

Batch no.	Strength	Type of Calcium hydrogenphosphate anhydrous	pH of 20% (m/V) slurry of capsule content
(CC)	5.0 mg	Di CAFOS A	6.10
(CX)	5.0 mg	Di CAFOS A	9.59

5

**Batch (CA) has been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- gelatine capsules have been filled with this powder blend

15

**Batch (CB) has been manufactured as follows:**

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.

- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.

- HPMC capsules have been filled with this powder blend

23



WO 02/053134

23

PCT/NL01/00606

Batch (CC) has been manufactured as follows:

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- All excipients except magnesium stearate have been mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.
- Magnesium stearate was added and the powder blend was mixed for another 5 minutes at about 25 rpm.
- *gelatine* capsules have been filled with this powder blend

10

Batch (CX) has been manufactured as follows:

- The amlodipine maleate was sieved through a 500 µm screen.
- The other excipients have been sieved through a 850 µm screen.
- Amlodipine maleate, magnesium oxide and about 30 % of the amount of microcrystalline cellulose (MCC) were mixed in a free fall mixer for 10 minutes at about 25 rpm.
- The remaining amount of MCC, calcium hydrogenphosphate anhydrous and sodium starch glycolate were added and the blend was mixed in a free fall mixer for 15 minutes at about 25 rpm. pH value was checked in 20% aqueous slurry.
- Magnesium stearate was added and the powderblend was mixed for another 5 minutes at about 25 rpm.
- *gelatine* capsules have been filled with this powderblend, using an automatic capsule filling machine.

25

#### Example 7 Stability studies on amlodipine maleate capsules

Stability studies on batches prepared in Example 6 were performed essentially as described in Example 5.

WO 02/053134

24

PCT/NL01/00606

Stability studies performed in PVC/PE/PVDC blister at 40 °C/75 % RH

t=0 months	(CA)	(CB)
Assay	5.13	4.98
Z#202	0.04	0.04
Z#203	0.03	0.03
Z#204	0.12	0.12
Z#205	0.00	0.00
tot unkn.	0.32	0.31
t=3 months		
Assay	4.82	4.76
Z#202	0.08	0.06
Z#203	0.02	0.02
Z#204	0.15	0.14
Z#205	0.01	0.00
tot unkn.	0.39	0.38
t=6 months		
Assay	4.67	4.81
Z#202	0.14	0.10
Z#203	0.04	0.02
Z#204	0.23	0.18
Z#205	0.02	0.01
tot unkn.	0.45	0.42

WO 02/053134

25

PCT/NL01/00606

Two one month stability studies performed in open dish

t=0 months	(CC)	(CX)
Assay	4.91	4.72
Z#202	0.04	0.04
Z#203	0.03	0.04
Z#204	0.12	0.13
Z#205	0.00	0.00
tot unkn.	0.28	0.29
T = 1 month @ 25°C/60%RH		
Assay	4.85	4.70
Z#202	0.05	0.05
Z#203	0.04	0.04
Z#204	0.13	0.14
Z#205	0.00	0.00
tot unkn.	0.28	0.28
T=1 month @ 40°C/75%RH		
Assay	4.76	4.16
Z#202	0.06	0.08
Z#203	0.04	0.03
Z#204	0.15	11.36
Z#205	0.00	0.00
tot unkn.	0.28	0.62

WO 02/053134

26

PCT/NL01/00606

Stability studies performed in HDPE container

t=0 months	(CA)	(CB)
Assay	4.99	5.01
Z#202	0.05	0.04
Z#203	0.03	0.03
Z#204	0.12	0.11
Z#205	0.00	0.00
tot unkn.	0.31	0.30
T= 3 months @ 25°C/60%RH		
Assay	5.07	4.77
Z#202	0.04	0.04
Z#203	0.03	0.04
Z#204	0.12	0.12
Z#205	0.00	0.00
tot unkn.	0.30	0.32
T= 3 months @ 40°C/75%RH		
Assay	5.01	4.63
Z#202	0.05	0.05
Z#203	0.03	0.04
Z#204	0.12	0.13
Z#205	0.00	0.00
tot unkn.	0.29	0.32

Stability studies on commercially available Amlor® (amlodipine besylate)

5 capsules

Stability studies performed in original blister

Amlor 5 mg capsules batch 9037002	t=0 months	t=3 months 40°C/75%RH
Assay	4.59	4.44
Z#202	0.01	0.20
Z#203	0.00	0.00
Z#204	0.00	0.00
Z#205	0.00	0.00
tot unkn.	0.06	0.37

WO 02/053134

PCT/NL01/00606

27

The invention having been described, it will be readily apparent to those skilled in the art that further changes and modifications in actual implementation of the concepts and embodiments described herein can easily be made or may be learned  
5 by practice of the invention, without departing from the spirit and scope of the invention as defined by the following claims

WO 02/053134

28

PCT/NL01/00606

**CLAIMS**

- 5           1.    A pharmaceutical composition comprising an effective amount of  
amlodipine maleate and at least one pharmaceutically acceptable excipient wherein  
said composition has a pH within the range of 5.5- 7.0.
2.    The composition according to claim 1, wherein said composition has a  
10   pH of about 6.0 - 7.0.
3.    The composition according to claims 1 or 2, wherein said composition  
is in solid form.
- 15          4.    The composition according to any of the preceding claims, wherein  
said excipient is calcium phosphate or microcrystalline cellulose.
5.    The composition according to claim 4, wherein said composition  
comprises calcium phosphate and microcrystalline cellulose.
- 20          6.    The composition according to claim 4, wherein said excipient is  
calcium hydrogen phosphate.
7.    The composition according to claim 4, wherein said excipient is  
25   microcrystalline cellulose.
8.    The composition according to any of the preceding claims, further  
comprising an acidic pH adjusting agent.
- 30          9.    The composition according to any of the preceding claims, wherein  
said composition is in the form of a tablet.

WO 02/053134

PCT/NL01/00606

29

10. The composition according to claim 9, which further comprises an outer moisture and/or light barrier layer surrounding said tablet.

11. The composition according to any of the preceding claims 1-8, wherein  
5 said composition is in the form of a capsule,

12. The composition according to any of the preceding claims, wherein said amount of amlodipine maleate corresponds to 1.0 to 25 mg of amlodipine free base.

10

13. The composition according to claim 12, wherein said amount of amlodipine maleate corresponds to 1.25, 2.5, 5 or 10 mg of amlodipine free base.

14. A method for treating or preventing angina or hypertension or heart  
15 failure, which comprises administering to a patient in need thereof an effective amount of the composition according to any of the claims 1-13.

15. A process for making the composition according to any of the claims 1, 3-13 or 2-13, which comprises mixing amlodipine maleate and at least one  
20 pharmaceutically acceptable excipient to form a mixture having a pH within the range of 5.5 to 7, or 6.0 - 7.0 respectively.

16. A process, which comprises:  
mixing amlodipine maleate and at least one pharmaceutically acceptable excipient to  
25 form a mixture having a pH of 5.5-7.0.

17. The process according to claim 16, which further comprises compressing said mixture into a tablet.

30 18. The process according to claim 16, which further comprises filling capsules with said mixture to form a pharmaceutical dosage form.

19. The process according to claim 16, wherein said mixing is carried out by wet granulation.

WO 02/053134

PCT/NL01/00606

30

20. The process according to claim 16, wherein said mixing is carried out by a dry method.

5 21. The process according to claim 20, wherein said amlodipine maleate is mixed as solid particles having an average particle size of at least 100 microns with said excipient.

22. A tablet made according to the process of any of the claims 16, 17, 19,  
10 20, 21.

23. A process, which comprises:  
mixing solid particles of amlodipine maleate, having an average particle size of at  
least 20 microns, with a pharmaceutically acceptable excipient to form a mixture.

15

24. The process according to claim 23, which further comprises filling  
capsules with said mixture to form a pharmaceutical dosage form.

25. The process according to claim 23, which further comprises  
20 compressing said mixture to form a tablet.

26. The process according to claim 23, wherein said average particle size is  
at least 100 microns.

25 27. The process according to claim 23, wherein said mixture is blended  
with one or more appropriate excipients so as to have a pH within the range of 5.5 to  
7.

28. The use of a composition according to any of the claims 1-13 for the  
30 manufacture of a medicament for the treatment of angina, hypertension or heart  
failure.



## INTERNATIONAL SEARCH REPORT

PCT/NL 01/00606

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 A61K9/20 A61K9/48 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 34299 A (TELOR OPHTHALMIC PHARMACEUTICALS) 21 December 1995 (1995-12-21) claims 11,14,15 page 7, line 27 - page 8, line 3	1,2,4, 13,15,16
A	WO 95 05822 A (PFIZER) 2 March 1995 (1995-03-02) the whole document	1-28
E	WO 01 74390 A (PFIZER) 11 October 2001 (2001-10-11) claims 1,3,9 page 4, line 1 - line 7 page 6, line 1 - line 15 page 7, line 4 - line 15 example 1	1-7,9, 11-20,28

— / —

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document relating to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application published to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

21 May 2002

Date of mailing of the international search report

29/05/2002

Name and mailing address of the ISA

European Patent Office, P.O. Box 1  
 NL - 2200 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tlx 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Ventura.Amat, A

# INTERNATIONAL SEARCH REPORT

PCT/NL 01/00606

C. (CONTINUATION) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	<p>EP 1 181 932 A (PFIZER)  27 February 2002 (2002-02-27)  claims 1,16  column 4, line 34 - line 42  column 5, line 8 - line 15  column 5, line 29 - line 42  example 4</p>	<p>1-4,7,9,  11-20,28</p>

# INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/NL 01/00606

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9534299	A	21-12-1995	WO 9534299 A1	21-12-1995
WO 9505822	A	02-03-1995	AT 192337 T	15-05-2000
			AU 686658 B2	12-02-1998
			AU 7612994 A	21-03-1995
			CA 2170278 A1	02-03-1995
			CN 1129907 A , B	28-08-1996
			DE 69424317 D1	08-06-2000
			DE 69424317 T2	24-08-2000
			DK 754043 T3	07-08-2000
			WO 9505822 A1	02-03-1995
			EP 0754043 A1	22-01-1997
			ES 2145149 T3	01-07-2000
			GR 3033547 T3	29-09-2000
			IL 110700 A	11-04-1999
			JP 2000044475 A	15-02-2000
			JP 3007992 B2	14-02-2000
			JP 8509004 T	24-09-1996
			KR 163840 B1	01-12-1998
			NO 960730 A	23-02-1996
			NZ 271993 A	24-11-1997
			PT 754043 T	29-09-2000
			US 6080761 A	27-06-2000
			ZA 9406475 A	26-02-1996
WO 0174390	A	11-10-2001	AU 3951001 A	15-10-2001
			WO 0174390 A2	11-10-2001
			US 2001036954 A1	01-11-2001
EP 1181932	A	27-02-2002	BR 0103434 A	26-03-2002
			EP 1181932 A2	27-02-2002
			US 2002045648 A1	18-04-2002